

Tuberculosis.—The next table sets out the death-rate from tuberculosis for 1940-1945.

Death-rates per 10,000 Population
(Europeans only)

Year.	Respiratory Tuberculosis.	Non-respiratory Tuberculosis.	Tuberculosis (all Forms).
1940	3·24	0·64	3·88
1941	3·19	0·69	3·88
1942	3·18	0·75	3·93
1943	3·09	0·63	3·72
1944	3·12	0·69	3·81
1945	3·12	0·66	3·78

Venereal Disease.—The following tables summarize the work of the venereal-disease clinics in the four main centres. Figures are given for the last pre-war years as well as for subsequent years. The reduction in the incidence of syphilis since 1941 has been maintained, and the figures now approximate the pre-war level :—

Number of Persons seen for the First Time at each Venereal-disease Clinic and found to be suffering from Syphilis

Year.	Auckland.		Wellington.		Christchurch.		Dunedin.		Total.		Grand Totals.
	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	
1939	59	43	18	25	19	11	12	5	168	84	192
1940	63	37	77	58	12	8	22	8	174	111	285
1941	102	57	96	63	29	17	33	6	260	144	403
1942	70	78	53	71	18	11	20	6	161	166	327
1943	48	95	20	41	17	14	29	3	114	153	267
1944	21	48	14	26	14	10	27	4	76	88	164
1945	61	34	11	20	15	8	27	6	114	68	182

Number of Persons seen for the First Time at each Venereal-disease Clinic and found to be suffering from Gonorrhoea

Year.	Auckland.		Wellington.		Christchurch.		Dunedin.		Total.		Grand Totals.
	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	
1939	399	141	333	53	328	88	91	90	1,151	372	1,523
1940	474	118	310	53	286	79	78	89	1,148	339	1,487
1941	410	183	373	42	271	72	81	79	1,135	376	1,511
1942	312	286	236	63	181	69	75	73	804	491	1,295
1943	265	441	138	89	122	92	51	15	576	637	1,213
1944	215	470	140	59	139	86	50	22	544	637	1,181
1945	389	413	178	54	149	66	46	9	762	542	1,304

Notifications of Disease.—Tables showing the notifications of infectious and other notifiable diseases are included in the section dealing with Public Hygiene.