$Complications{\rm--continued.}$

	Cen- tral.	Marg. with Hæm.	Without Hæm.	Remarks.
Placenta previa	No. 2	No. 9	No. 2	I central, at Auckland, child stillborn and with concealed hæmorrhage; I marginal, child born alive, prolapse of cord: rapid dilatation in both cases with Bozzi. Several of the other cases at Dunedin only slight hæmorrhage; none very severe.

Morbidity List (with Day of Occurrences of Temperature).

Temp. for at least	Num.	Days.											
Two Consecutive Days.	Total I	lst.	2nd.	3rd.	4th.	őth.	6th.	7th.	8th.	9th.	10th and	over.	Causes.
100 to 100·8	19	6	5	1		3		1		1	2		Adhered placentæ 2, fissured nipples, mastitis 4, lacerations 3, constipation, bronchitis, abscess of breast, syphilitic goitre infection, septicæmia 4, sapræmia 12, ex-ophthalmic goitre puerperal
100·8 to 102	18	3	2	2	2	7	1	1	• •			}	mania, acute rheumatism, influenza 3, pneu- monia, pyonephritis following eclampsia, phle- bitis, cystitis, metritis, scarlet fever 1, phthisis,
102 to 103	15	1	3	3	2	1	•••	2	1	2			degenerate chorion 1, phlegmasia 1, anæmia with albuminuria 3, albuminuria with head disease 1, abdominal tuberculosis and mastoid
Over 103	15	3	1	4	4	1	1	 			1	}	abscess 1, intestinal sepsis from foul teeth 2, pyæmia after old pelvic cellulitis 1.

Mortality.

		No.	Remarks.
Maternal	••	5	1 emergency case outside, eclampsia, Christchurch; I septicæmia, Christchurch; 1 septicæmia and phthisis, Auckland; 1 cardiac failure with albuminuria after twin labour, Wellington; 1 embolism of 12th day of normal confinement, Wellington.
Infantile	••	6	I overlain (out-patient); I congenital heart-disease, Auckland; I malæna neonatorum; I congenital malformation; I weakly from birth; I spasm of glottis, with inability to swallow.
Stillbirths, term	full	12	1 placenta prævia; 3 albuminuria; 1 second twin, albuminuria; 1 hydrocephalus; 1 transverse; 1 cardiac lesion in mother.
Premature		11	3 triplets; 3 twins; 1 albuminuria in mother.
Macerated		8	1 severe albuminuria, with toxic symptoms.

Inability to Nurse.

		No.	Remarks.
Complete	••	25	In 2 lactation was never established, elderly primiparæ; in 1 case child was feeble; 1 premature child left in hospital; 1 had eclampsia, followed by
Partial	••	68	pyelitis; 10 had anæmia; 1 septicæmia; 7 had phthisis; 1 mammary deformity; and 3 badly developed breasts and nipples. Of these cases, many had insufficient or poor supply; 5 had pyrenia; 2 plural pregnancies; 4 anæmia; and several had albumin with toxic symptoms.