

## COMPLICATIONS.

—	No.	Alb. pres. on Dis- charge.	Alb. abs. on Dis- charge.	Accompanying Complications.
Albuminuria before labour	104	No. 26	No. 74	1 case due to injury to kidney four days before labour; 1 had albuminuric retinitis; 5 adherent placenta; subinvolution, post-partum hæmorrhage, 6; retained placenta, 5; eclampsia, 2.
With toxic symptoms ..	16	1	1	1 secondary post-partum hæmorrhage; 1 child was anencephalic; one had bad talipes varus; 1 spina bifida with paralysis of lower legs.
Eclampsia .. ..	7	..	..	

—	Central.	Marg. with Hæm.	Without Hæm.
	No.	No.	No.
Placenta previa .. ..	1	8	2 lateral.

## MORBIDITY LIST.

(With Day of Occurrences of Temperature.)

Temp. for at least Two Consecutive Days.	Total No.	Days.										Some Causes.
		1st.	2nd.	3rd.	4th.	5th.	6th.	7th.	8th.	9th.	10th & over.	
100 to 100·8 ..	29	6	5	8	4	2	2	2	..	..	..	8 sapræmia; 10 albuminuria; 3 toxæmia; 1 influenza; 6 mastitis; 1 exophthalmic goitre; 1 cellulitis.
100·8 to 102 ..	19	9	1	1	3	..	..	1	..	2	1	2 sapræmia; 1 oval sepsis; 1 pneumonia; 1 influenza; 1 nephritis; 1 diabetes; 1 mental; 5 mastitis.
102 to 103 ..	10	7	2	..	2	..	1	..	..	..	1	2 inflamed varicose veins; 2 eclampsia; 1 influenza.
Over 103 ..	10	4	..	1	2	1	1	1	..	..	..	1 constipation; 1 influenza; 2 sapræmia; 1 scarlet fever; 1 cellulitis.

## MORTALITY.

—	No.	Remarks.
Maternal ..	2	1 primipara from eclampsia; 1 concealed hæmorrhage admitted moribund and died shortly after admission.
Infantile ..	17	1 patent foramen ovale; 1 anencephalic; 1 asphyxiated; 1 atelectasis; 1 cerebral hæmorrhage; 1 bronchitis and syncope; 1 malassimilation; 1 convulsions; 1 failure to establish circulation; 7 premature; 1 albuminuria in mother.
Still-births, full term	..	2 placenta prævia; 3 prolapse of cord; 1 deformed foetus; 1 cord many times round neck, not pulsating at birth.
Premature ..	11	6 under 7 months; 1 ante-partum hæmorrhage; and 1 eclampsia in mother.
Macerated ..	14	Mothers with albuminuria, eclampsia, &c.

## INABILITY TO NURSE.

—	No.	Remarks.
Complete ..	20	Due to anæmia; cicatrices due to burns; defective breasts; inverted nipples; heart and kidney trouble; acute diabetes; chronic nephritis; severe mastitis; 2 abscess of breast; consumption; general poor health; albuminuria; retracted nipples; prematurity.
Partial ..	113	Large number due to anæmia; supplementary feeds twice during day given; many able to nurse entirely at third week. Milk-and-water, humanized milk, milk and barley-water, and Glaxo were used.