

APPENDIX III.

NOMENCLATURE.

LIST SHOWING POPULAR AND SCIENTIFIC NAMES OF FISH, CRUSTACEA, AND MOLLUSCA MENTIONED IN REPORT.

FISHES—	Popular Names.	Scientific Names.
Barracouta		<i>Thyrssites atun</i> (Euphrasen).
Bass (or bass groper)		<i>Polyprion americanus</i> (Bloch. and Schn.).
Blue cod		<i>Parapercis colias</i> (Forster).
Brill		<i>Colistium ammotretis guntheri</i> (Hutton).
Butterfish (= greenbone or kelp-fish)		<i>Coridodax pullus</i> (Forster).
Conger		<i>Leptocephalus conger</i> (Linnaeus).
Dab (commonly included among flounder)		<i>Rhombosolea plebeia</i> (Richardson).
Elephant-fish		<i>Callorhynchus milii</i> (Bory).
Flounder		<i>Rhombosolea leporina</i> Guenther.
Frost fish		<i>Lepidopus caudatus</i> (Euphrasen).
Greenbone (= butterfish)		<i>Coridodax pullus</i> (Forster).
Groper (= hapuku)		<i>Polyprion oxygeneios</i> (Bloch. and Schn.).
Gurnard		<i>Chelidonichthys kumu</i> (Lesson and Garnot) and <i>Lepidotrigla brachyoptera</i> Hutton.
Hake (= southern kingfish)		<i>Jordanidia solandri</i> (Cuv. and Val.).
Hapuku (= hapuka or whapuku)		<i>Polyprion oxygeneios</i> (Bloch. and Schn.).
Herring (= South Island mullet)		<i>Agonostomus forsteri</i> (Cuv. and Val.).
“ Picton herring ” (= pilchard or sardine)		<i>Sardinia (Clupea) neopilcharda</i> (Steindachner).
John Dory (or dori)		<i>Zeus faber</i> Linnæus.
Kahawai		<i>Arripis trutta</i> (Forster).
Kingfish (northern)		<i>Seriola lalandi</i> (Cuv. and Val.).
Kingfish (southern)		<i>Jordanidia solandri</i> (Cuv. and Val.).
Ling		<i>Genypterus blacodes</i> (Bloch. and Schn.).
Maomao (maumau)		<i>Scorpius violaceus</i> (Hutton).
Moki		<i>Latridopsis ciliaris</i> (Forster).
Mullet (northern)		<i>Mugil ?cephalus</i> Linnæus.
Mullet (southern) (= herring)		<i>Agonostomus forsteri</i> (Cuv. and Val.).
Pioke (dog-fish)		<i>Squalus fernandinus</i> (Molina).
Piper (garfish)		<i>Hemirhamphus intermedius</i> Cantor.
Red cod		<i>Physiculus bachus</i> (Bloch. and Schn.).
Rock-cod		<i>Lotella rhacinus</i> (Forster).
Sardine		<i>Sardinia (Clupea) neopilcharda</i> (Steindachner).
Skate		<i>Raja nasuta</i> Mueller and Henle.
Snapper		<i>Pagrosomus auratus</i> (Forster).
Sole		<i>Peltorhamphus novae-zeelandiae</i> Guenther.
Tarakihi		<i>Dactylopagrus macropterus</i> (Forster).
Trevally		<i>Caranx platessa</i> (Cuv. and Val.).
Trumpeter		<i>Latris lineata</i> (Forster).
Turbot		<i>Colisteum nudipinnis</i> (Waite).
Warehou		<i>Seriola brama</i> (Guenther).
Whitebait		<i>Galaxias attenuatus</i> (Jenyns).
CRUSTACEA—		
Crayfish		<i>Jasus lalandii</i> (Milne-Edwards) and <i>J. hugeli</i> (Heller).
Whale-feed		<i>Munida gregaria</i> Fabricius.
MOLLUSCA—		
Borer		<i>Thais scobina</i> Quoy and Gaimard.
Cockle (or Pipi)		<i>Chione (Antigona) stutchburyi</i> Gray.
Mussel		<i>Mytilus canaliculus</i> Martyn.
Oyster, dredge		<i>Ostrea sinuata</i> Lamarek (<i>O. angasi</i> Sowerby).
Oyster, rock		<i>O. glomerata</i> Gould.
Pipi		<i>Chione, Dosinia, Mesodesma subtriangulatum</i> , &c.
Pupu		<i>Thais succincta</i> Martyn.
Toheroa		<i>Mesodesma (Amphidesma) ventricosa</i> Gray.