

Table VI—continued.

Group.	Reason given for Operation and Parity.	Number of Cases.	Number of Deaths.		Cause of Deaths of Mothers, and Notes on Special Cases.
			Infants.	Mothers.	
III	Placenta prævia—				
	1 para	5	4	..	..
	2 para	3	2	..	..
	3 para	2	..	..	..
	5 para	1	..	..	..
	6 para	1	1	..	..
	13 para	1	1	..	..
	Not stated	1	1	..	..
	Total	14	9	..	
IV	Accidental hæmorrhage—				
	1 para	3	3	1	Hæmorrhage: age thirty-two, three miscarriages in four years, all due to toxæmia.
	2 para	1	..	..	..
	4 para	1	1	..	..
	6 para	1	1	..	..
	Not stated	2	2	1	Hæmorrhage.
	Total	8	7	2	
V	Eclampsia—				
	1 para	5	..	..	..
	Not stated	2	2	1	Pulmonary embolism and ante-partum hæmorrhage.
	Total	7	2	1	
VI	Toxæmia of pregnancy—				
	1 para	12	2	1	Sub-acute nephritis.
	2 para	1	1	..	..
	4 para	1	1	1	Pulmonary embolism: six-month pregnancy.
	Not stated	2	..	..	..
	Total	16	4	2	
VII	Other conditions—				
	Heart-disease—				
	1 para	1	..	..	..
	2 para	1	..	..	..
	Right lobar pneumonia	1	..	..	..
	Detachment of retina, danger of blindness—				
	2 para	1	..	..	..
	Hydronephrosis—				
	1 para	1	1	..	..
	Bartholin's cyst—				
	1 para	1	1	..	..
	Secondary carcinoma of pelvis—				
	1 para	1	1	..	..
	Vasco-vaginal fistula—				
	3 para	1	..	..	..
	Weak uterine: Scars due to previous operations—				
	1 para	1	..	..	Uterus weakened by previous removal of fibroids.
	2 para	2	..	..	..
	6 para	1	1	..	Uterus ruptured through old cæsa-rean scar.
	Total	12	4	..	
	Total cases	157	34	12	

Case-mortality rate, 7·64 per cent. Infant-mortality rate, 21·65 per cent.