H.—31.

75

Of the total of 1,018 cases notified during the year, records are available of the age-grouping of 852, of which 515 were in Auckland District and 377 in Wellington District.

Table 2.—Showing Distribution by Age-Groups, of 852 Cases of Poliomyellitis, 1916.

	-				Auckland District.	Wellington District.
Under 5 years			 		322	200
5 to 10 years			 		84	65
10 to 15 years			 		43	27
15 to 20 years			 		35	19
20 to 30 years			 		23	18
30 to 40 years			 		5	7
40 years and over			 	• •	3	1
		Totals	 		515	337

Table 3.—Showing Deaths, by Age-Groups, from Poliomyelitis among 852 Cases, 1916.

	 _	Auckland District.	Wellington District.		
Under 5 years	 	 		23	20
5 to 10 years	 	 		11	3
10 to 15 years	 	 		5	4
15 to 20 years	 	 		8	9
20 years and over	 	 		6	10
	Totals	 		53	46

Table 4.—Acute Poliomyelitis, 1924–25: Table showing Notifications, by Health Districts and Week of Notification.

Week ended		North Auckland.	Central Auckland.	South Auckland.	Coromandel- Opotiki.	Taranaki- Horowhenua.	Wairarapa- East Cape.	Central Wellingcon.	Nelson- Marlborough.	West Coast.	Canterbury.	Otago.	Southland.	Totals.
24th November, 1924 1st December, 1924 8th December, 1924 15th December, 1924 22nd December, 1924 22nd December, 1924 29th December, 1924 25th January, 1925 12th January, 1925 26th January, 1925 26th January, 1925 26th February, 1925 27th February, 1925 27th March, 1925 27th March, 1925 23rd March, 1925 23rd March, 1925 25th April, 1925		N 9 10 10 10 10 10 10 10 10 10 10 10 10 10	2 2 3 3	1 1 4 5 2 2 4 7 3 3 6 6 6 1 1 5 5 3 2 2 2 3	6do	20 21 22 24 47 26 30 21 21 22 24 47 26 21 21 22 24 47 47 47 47 44 43 44 47	Nai	3 6 11 9 9 17 17 26 14 10 4 1 6 3 2 2			1			1 3 3 6 6 13 11 12 19 300 600 600 600 138 88 104 1388 118 70 109 882 48 39 92 28 8 19 17
11th May, 1925 18th May, 1925 25th May, 1925 1st June, 1925 15th June, 1925 22nd June, 1925 29th June, 1925	 	I 1	1	1 1 1 1 		3 1 3	1 2				4 4 2 2 3 1 1	3 2 7 5 2 4	 1 1	13 13 14 7 6 5

The year 1917 showed an incidence above the average, but the disease remained sporadic until 1920, when 76 cases occurred, widely distributed, 28 of the cases being in Auckland. In 1921 a minor epidemic was experienced. Of the 267 cases notified, the North Island, with approximately double the population of the South Island, was responsible for 132. The increased incidence the following year was mainly in the North Island. Only a few cases were notified between 1922 and December, 1924, when the second major epidemic commenced.